

The Recent Results of Forward Regime Studies by the BRAHMS Experiment at RHIC

Natalia Katryńska*

for the BRAHMS Collaboration†

*Smoluchowski Institute of Physics,
Jagiellonian University*

4 Reymonta St., Kraków, Poland

(Received November 1st, 2009)

With regard for BRAHMS experiment unique opportunity to measure particle production in the wide range of rapidity the latest results of properties of the medium evolution from $y=0$ to the forward region in the ultra-relativistic collisions are presented. The research of the net-baryon density issue is shown for nucleus+nucleus system. The studies of baryon enhancement to the pion yield at intermediate p_T are presented for p+p and Au+Au reactions. Moreover, the BRAHMS measurements provide a great test of Color Glass Condensate (CGC) state by studying the nuclear modification factor, R_{dAu} , vs. rapidity for charged hadrons.

PACS numbers: 25.75.-q , 13.85.-t , 25.75.Dw

Keywords: forward rapidity, nuclear stopping, particle ratios, nuclear modification factor

I. INTRODUCTION

The BRAHMS experiment at RHIC can measure and identify particle over the wide range of rapidity and transverse momentum. It allows to study the different aspects of strong interactions in the ultra-relativistic nucleus-nucleus collisions. BRAHMS has estimated the degree of nuclear stopping by measurement of net-baryon distribution which is related to baryon number transport during the collision [1, 2]. The rapidity dependent proton-to-pion ratio reveal which picture of hadronization is supported in the bulk medium [3, 4]. The nucleon+nucleus data are a starting point of exploration of the initial state effects [5, 6]. In this paper, the recent BRAHMS results are presented for d+Au reactions at $\sqrt{s_{NN}} = 200$ GeV and for Au+Au and p+p collisions at $\sqrt{s_{NN}} = 62.4$ and 200 GeV. The description of BRAHMS experimental setup can be found in [7].

II. NET-BARYON DENSITY DISTRIBUTION

Wide rapidity acceptance of the BRAHMS

spectrometers provides unique opportunity to study nuclear stopping in the ultra-relativistic nucleus-nucleus reactions. The average rapidity loss, $\langle\delta y\rangle = y_b - \langle y\rangle$, quantifies stopping in heavy ions collisions. If initial baryon participants loose all the kinetic energy ($\langle\delta y\rangle = y_b$) we observe full stopping, for $\langle\delta y\rangle = 0$ the system is completely transparent. Before RHIC era, at Schwer Ionen Synchrotron (SIS) in Darmstadt, Alternating Gradient Synchrotron (AGS) in Brookhaven National Laboratory and Super Proton Synchrotron (SPS) in CERN it was noticed that $\langle\delta y\rangle$ is proportional to the beam rapidity [8]. The BRAHMS results [9] show that this linear scaling is broken above top SPS energies - Figure 1. Assuming that underlying physics is the same at highest attainable energies the extrapolation of $\langle\delta y\rangle$ to the beam rapidity characteristic for Large Hadron Collider is depicted in Figure 1.

While the nuclei are colliding, the nucleons, at first bounded up inside nuclei, loose their kinetic energy. The energy loss per participant baryon is estimated that $\langle\delta E\rangle = 21 \pm 2$ GeV for Au+Au @ $\sqrt{s_{NN}} = 62.4$ GeV [9] and $\langle\delta E\rangle = 73 \pm 6$ GeV for Au+Au @ $\sqrt{s_{NN}} = 200$ GeV [10] for the most (0-10%) central collisions. For two colliding nucleons the calculated energy loss constitutes 70% of the initial energy of beam. This energy loss is transformed mainly into particle production and random (thermal) motion of produced quanta.

*n.katrynska@if.uj.edu.pl

†The full name list of the BRAHMS Collaboration can be found at <http://www4.rcf.bnl.gov/brahms/WWW>




FIG. 1: The rapidity loss as a function of beam rapidity for heavy ions collisions at AGS, SPS and RHIC. The solid line represents the fit to the SPS and RHIC data. The grey band is the statistical uncertainty. The dashed line is linear fit to the AGS and SPS results [8]. Published in [9].

From BRAHMS measurements at the top RHIC energy one can conclude that the midrapidity region of the collision is almost net-baryon-free. It corresponds to picture of interacting matter proposed by Bjorken [1] with near free net-baryon content at midrapidity. At lower energy, $\sqrt{s_{NN}} = 62.4$ GeV, for Au+Au reactions at $y \approx 0$ the net-proton $\frac{dN}{dy}$ indicates that the medium is also quite transparent compared with data at SPS and AGS energy [8].

III. PROTON-TO-PION RATIO AT INTERMEDIATE P_T


The proton-to-pion ratios for Au+Au reactions at $\sqrt{s_{NN}} = 62.4$ GeV and $\sqrt{s_{NN}} = 200$ GeV give the possibility of capture the properties of bulk medium by studying baryon and meson production. As it has been shown in [11] for positively charged hadrons for heavy ion reactions the enhancement of baryon production in respect to the meson yield has been noticed at intermediate p_T . At the top RHIC energy of colliding nuclei the proton and pion production seems to be comparable with each other at midrapidity at

$2.5 \text{ GeV}/c < p_T < 4 \text{ GeV}/c$, but with increasing η the baryon production prevails over meson production successively. The centrality dependence of proton-to-pion ratio suggests that the p/π^+ ratios at intermediate transverse momenta scales with the size of the created medium. The ratio of \bar{p}/π^- reaches the maximum of 0.5 at $\eta \approx 2.25$ for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and decreases for more forward pseudorapidities.

The p/π^+ ratio for Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV for $\eta \approx 2.67, 3.2, 3.5$ is depicted in Figure 2. Additionally, the p+p results at the same energy and η bins are shown. At all pseudorapidities unexpected high value of 10 at $p_T = 1.5$ GeV/c of proton-to-pion ratio is observed [12]. There is remarkably little difference in the p/π^+ ratios from a very wide range of systems. This is in contrast to the trends at midrapidity and forward rapidity regimes for Au+Au at $\sqrt{s_{NN}} = 200$ GeV where significant medium effect reflected in dependence of p/π^+ ratios on system size is observed [11]. However, the observed consistency between the results of p+p reactions and Au+Au system for all centrality bins for $\eta \approx 3.2$ is reckoned as the crossing point in pseudorapidity. These results indicate that the nuclear modification factor for protons and pions are equal in the observed p_T range. It can be seen from:


$$\begin{aligned} R_{AA}^{\text{protons}} &= \frac{d^2 N_{\text{protons}}^{A+A}/dp_T dy}{d^2 N_{\text{pions}}^{A+A}/dp_T dy} = \\ R_{AA}^{\text{pions}} &= \frac{d^2 N_{\text{protons}}^{p+p}/dp_T dy}{d^2 N_{\text{pions}}^{p+p}/dp_T dy} = \\ &= \frac{p/\pi^+ (\text{Au} + \text{Au})}{p/\pi^+ (p + p)} = 1. \end{aligned} \quad (1)$$

According to studies of the net-baryon distribution BRAHMS data show rather transparent region at central rapidities with baryons shifted by two units from beam rapidity. This process intensifies longitudinal energy dissipation and contributes through coalescence mechanism to increased proton yield at intermediate p_T . At very forward rapidities one can expect less power of nuclear stopping which might be observed as a result of enhancement of proton production to pion production that, particularly, is seen for elementary reaction. The higher value of p/π^+ for $\eta \approx 3.5$ for p+p collisions might acknowledge this


 FIG. 2: The p/π^+ ratio vs. transverse momentum for Au+Au and p+p collisions at $\sqrt{s_{NN}} = 62.4$ GeV.

argumentation. The crossing point was predicted by UrQMD [13], HIJING [14] and AMPT [15] calculations, but almost one unit of rapidity lower as experimental data indicate.

Figure 3 shows the p/π^+ ratios as a function of p_T in the pseudorapidity range $2.6 \leq \eta \leq 3.8$ extracted from p+p reactions at $\sqrt{s_{NN}} = 200$ GeV. The behaviour of $p/\pi^+(p_T)$ ratio for p+p system reveals the characteristic tendency, namely the increase of peak value with increasing pseudorapidity. A very clear difference is found as the pseudorapidity changes from $\eta = 2.6$ to $\eta = 3.8$. The value of ratio grows systematically with rising η from 0.4 at $\eta = 2.3$ reaching almost 1 at the most forward pseudorapidity interval, $\eta \approx 3.8$. That high value of p/π^+ ratio is indispensable related to the large proton yield at high p_T . If we look carefully, especially for $\eta = 2.6, 3.1, 3.3$ the proton-to-pion ratio at highest covered p_T seems to converge a common value of 0.4 which is consistent with perturbative QCD predictions [3]. For negatively charged hadrons, the value of proton-to-pion ratio for elementary reactions in covered pseudorapidity interval reaches maximum value of 0.2 - Figure 4. The p/π^+ ratio exceeds the \bar{p}/π^- by a factor of 5. The peak value is less than 0.3 for $\eta = 2.3$ and dropping to < 0.1 for $\eta = 3.8$. Moreover, the maximum of \bar{p}/π^- ratio is shifted to the lower transverse momentum ($p_T \approx 1$ GeV/c) compared with the


 FIG. 3: The p/π^+ ratio for set of η bins in the range $2.6 \leq \eta \leq 3.8$ for p+p reactions at $\sqrt{s_{NN}} = 200$ GeV.

positive particles.

The interesting picture of \bar{p}/π^- ratios is revealed at low p_T . On the right hand of Figure 4 the comparison of \bar{p}/π^- for heavy ion and elementary reactions at $\sqrt{s_{NN}} = 200$ GeV is shown. The experimental data at the covered pseudorapidity interval, $2.6 \leq \eta \leq 3.8$, expose the same feature and the value of $\eta \approx 3.3$ is chosen just for clarity of the picture. Below $p_T \approx 0.9$ GeV/c




FIG. 4: The \bar{p}/π^- ratio vs. transverse momentum for $2.6 \leq \eta \leq 3.8$ for p+p collisions at $\sqrt{s_{NN}} = 200$ GeV (left hand). The results of ratio for p+p reactions are compared with data for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV for two intervals of centrality 0-10% and 40-80% at the same value of pseudorapidity, $\eta \approx 3.3$ (right hand).

the \bar{p}/π^- ratio increases towards more peripheral reactions for Au+Au collisions and reaches the largest value for p+p reactions. Moreover, the ratios for Au+Au for different centrality classes and for p+p system cross each other at approximately the same point, located $p_T \approx 0.9$ GeV/c. The higher \bar{p}/π^- for elementary system than for Au+Au collisions for soft p_T region might be the evidence of medium effects in heavy ions collisions at relativistic energies.

IV. RAPIDITY DEPENDENT NUCLEAR MODIFICATION FACTOR FOR D+AU COLLISIONS

Comparing the nuclear modification factor for pions, kaons and protons at $y = 0$ for d+Au reactions at $\sqrt{s_{NN}} = 200$ GeV we can observe a significant enhancement, especially for protons near $p_T \approx 2$ GeV/c [16]. This enhancement is known from SPS measurements [17] as the Cronin effect and is associated with partonic multiple scattering - an initial state broadening of the distribution of quark momenta. When we go to more forward rapidities BRAHMS has observed a significant suppression at high p_T . In Figure 5 the results of nuclear modification factor for identified hadrons for 0-30% centrality at $y \approx 3$ for d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV are presented. For

central collisions at forward rapidity the suppression of R_{dAu} is evident for all kinds of hadrons and there is no noticeable difference between individual species. The existence of the Color Glass Condensate might be an explanation of this effect in forward regime [5] where the low- x components (mostly gluons) of the wave function of the gold nuclei are probed by deuteron parton rescattered into forward region. On the other hand, in Dual Parton Model the addition of dynamical shadowing correction has been incorporated to describe the R_{dAu} behaviour [6].

V. SUMMARY

The net-baryon density results show the linear scalability of rapidity loss at SPS energies is broken at RHIC and slowly depends on beam rapidity that might be interpreted in the context of gluon saturation. The wide rapidity range studies of the $p/\pi(p_T)$ indicate how the hadronization process is driven with changing bulk medium properties. For elementary reactions the rapidity dependence of ratios is significant for both charges of hadrons. The data of p/π^- ratio for p+p collisions at $\sqrt{s_{NN}} = 200$ GeV rise above the results for heavy ion reactions for forward η at low p_T . At lower collision energy the crossing point in pseudorapidity is found for p+p and all




FIG. 5: The nuclear modification factor R_{dAu} of π^+ , K^+ and p for d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV at $\eta \approx 3$. The R_{dAu} of negatively charged hadrons marked with blue (online) squares was published in [16].

centrality intervals for Au+Au system. The studies of R_{dAu} for identified hadrons at $\sqrt{s_{NN}} = 200$ GeV are the foreword of existence of the Color

Glass Condensate state, however new evidences are coveted.

-
- [1] J. D. Bjorken, Phys. Rev. D **27**, 140-151 (1983).
 [2] L. D. Landau, Izv. Akad. Nauk SSR **17**, 51 (1953).
 [3] T. Hirano, Y. Nara, Nucl.Phys. A **743**, 305 (2004).
 [4] R. Hwa, J. Phys. G: Nucl. Part. Phys. **35**, 104017 (2008).
 [5] D. Kharzeev, Y. Kovchegov, K. Tuchin, Phys. Rev. D **68**, 094013 (2003).
 [6] A. Capella, E. G. Ferreiro, A. B. Kaidalov, D. Sousa, arXiv:0403081.
 [7] BRAHMS Collab., Nucl. Instr. Meth. A **499**, 437 (2003).
 [8] F. Videbæk, O. Hansen, Phys. Rev. C **52**, 2684 (1995).
 C. Blume (NA49 Collab.), J. Phys. G **34**, 951-954 (2007).
 [9] BRAHMS Collab., Phys. Lett. B **677**, 267-271 (2009).
 [10] BRAHMS Collab., Phys. Rev. Lett. **93**, 102301 (2004).
 [11] BRAHMS Collab., arXiv:0910.3328.
 [12] R. Hwa, C. B. Yang, Phys. Rev. C **76**, 014901 (2007).
 [13] <http://th.physik.uni-frankfurt.de/ürqmd>, July 2009.
 [14] <http://www-nsdth.lbl.gov/~xnwang/hijing>, July 2009.
 [15] Z. W. Lin, C. M. Ko, S. Pal, Phys. Rev. Lett. **89**, 152301 (2002).
 [16] E.-J. Kim, H. Yang for BRAHMS Collab., INPC2007, (2007).
 [17] J. W. Cronin, H. J. Frisch, M. J. Shochet, Phys. Rev. D **19**, 764 (1979).