DM-TPC: a new approach to directional detection of Dark Matter

Gabriella Sciolla

MIT

Outline:

- Introduction to Dark Matter
- Why directional detection of DM
- DM-TPC: detector concept
- Recent results: first evidence of "head-tail" effect
- Next step: toward a full-scale detector
- Conclusion

BNL Nuclear Physics Seminar, January 2008

First hints of Dark Matter

Fritz Zwicky (1933)

- Applying virial theorem to study of Coma cluster, he concluded that mass of galaxies in cluster was $O(10^2)$ what inferred from luminosity
- Consequence: substantial amount of matter not emitting light (Dark) must exist

Coma Cluster in ultraviolet and visible light from Sloan Digital Sky Survey/Spitzer Space Telescope

Strong evidence for DM

Vera Rubin et al. (1979)

- Study of rotational curve of spiral galaxies
  - Newtonian prediction for orbital velocity of galaxies

$$\frac{GMm}{r^2} = \frac{mv^2}{r} \Rightarrow \boxed{v \propto \frac{1}{\sqrt{r}}}$$

  - Observation: orbital velocity is flat outside central bulge
- Explanation: substantial amount of matter far from the center of the galaxy that is not emitting light (Dark Matter)



Even more convincing evidence…

Bullet Cluster (2006)

- Two colliding clusters of galaxies
- Its components (stars, gas, and DM) behave differently during collision
  - Stars (optical) not greatly affected: small gravitational slow down
  - Hot gas (X-rays), larger mass, EM interactions: more dramatic slow down
  - DM (gravitational lensing), largest mass, minimally affected
- Conclusion: most of the mass in the cluster pair is in the form of weakly interacting Dark Matter

X-ray: NASA/CXC/CfA/ M.Markevitch et al.;
Lensing Map: NASA/STScI; ESO WFI; Magellan/U.Arizona/ D.Clowe et al
Optical: NASA/STScI; Magellan/U.Arizona/D.Clowe et al.;

What is Dark Matter made of?

- Astrophysics and cosmology tell us that Dark Matter accounts for a huge fraction of our Universe:
  - 85-90% of the mass
  - 24% of the energy



- What is Dark Matter?
  - Many candidates:
    - Baryonic DM (e.g.: non-luminous gas)
    - Non baryonic DM --- hot or cold
  - CMB data favor cold non-baryonic Dark Matter
    - Cold: large mass --> non-relativistic velocities
    - Non-baryonic: gravity and weak interactions --> new particle
    - Stable: maybe LSP?
- Weakly Interacting Massive Particles (WIMPs) are the most likely candidates

Direct detection of WIMPs



Signal:

$\chi N \rightarrow \chi N'$

Backgrounds:

$n\ N \rightarrow n\ N'$

$\gamma\ e^- \rightarrow \gamma\ e^{-\prime}$

$N \rightarrow N' + \alpha, e^-$

$\nu\ N \rightarrow \nu\ N'$

- Basic principle: detect recoil of matter after elastic scatter with WIMP
- Different experiments use different techniques and materials
  - Ionization, scintillation, phonons
  - Si, Ge, CsI, Xe, Ar, CF_4, …
- Challenging measurements:
  - Very low-energy recoils (10-100 keV), very weak interactions, many backgrounds

Present DM searches



Current best limits on spin-independent interactions from:

___ Xenon-10
___ CDMS
___ ZEPLIN
___ WARP

- Many experiments engaged in direct detection of DM
  - Recent progress: improved cross-section limits $\sigma_{SI} < 10^{-44}$-10^{-43} cm^2
- Intrinsic limitation of mainstream DM experiments:
  - Counting experiments: zero-background assumed
  - Larger detectors will start to see several (irreducible) backgrounds

It will be very hard for a counting experiment
to provide unambiguous positive observation of Dark Matter

Situation similar to neutrino oscillations…

Ray Davis



First SNO results



Homestake experiment



Oscillation of solar ν first observed by Davis in the 60's in counting experiment, but decisive proof came in 2001 with water Cherenkov directional results

Dark Matter wind from Cygnus

The decisive proof of positive observation of DM requires correlation with astrophysical phenomena



Animation by A-M Piso

A wind of Dark Matter from Cygnus

The decisive proof of positive observation of DM requires correlation with astrophysical phenomena



Why not yearly asymmetry?



Yearly asymmetry:

- Small rate asymmetry: 2-10%
- Hard to disentangle from temperature dependent phenomena

Drukier, Freese, Spergel (1987)

Unambiguous signature of Dark Matter

Daily asymmetry~30-100%!



Only directional detection can correlate with Cygnus:
unambiguous positive observation of Dark Matter in presence of backgrounds

Directional Detectors

- Direction of incoming WIMP is encoded in direction of nuclear recoil





- How to detect the direction of recoils?
  - Low-pressure gaseous detectors
    - A 50 keV F in CF_4 @ 40 torr recoils ~2 mm

Spin-dependent interactions

- WIMPs can scatter elastically on nuclei via
  - Spin-independet interactions
    - cross-section scales with the mass of the nucleus: $\sigma \sim A^2$
  - Spin-dependent interactions
    - cross-section is nonzero only if the nucleus has a nonzero spin
- Spin-dependent interactions may be enhanced by orders of magnitude compared to spin-independent
  - E.g.: in models in which LSP has substantial Higgsino contribution

Chattopadhyay and D.P. Roy, Phys. Rev. D 68(2003) 33010
Murakami B. and J.D. Wells, Phys. Rev. D 64 (2001) 15001
Vergados, J., J. Phys. G 30 (2004) 1127

- Weaker limits for spin-dependent interactions
  - Limits on spin-independent x-section: $\sim 10^{-44}$-10^{-43} cm^2
  - Limits on spin-dependent x-section: $\sim 10^{-37}$-10^{-36} cm^2

7 orders of magnitude!

Spin-dependent searches are promising and almost unexplored

Our goal

Develop a novel detector for direct detection of Dark Matter with the following characteristics:

- Directionality
  - The only way to provide unambiguous positive observation of Dark Matter in presence of backgrounds
- Spin-dependent interactions
  - Can be much enhanced wrt spin-independent interactions

To make this feasible we need:

- Low cost/unit volume
  - Directionality requires gaseous detectors: large volumes
- Easy to maintain
  - Very stable, safe, easy to operate underground
- Scalability
  - Modular structure

The DM-TPC Collaboration

S. Ahlen, D. Avery*, H. Tomita, A. Roccaro, M. Lewandowska[1]
Boston University

N. Skvorodnev, H. Wellenstein
Brandeis University

B. Cornell*[2], D. Dujmic, W. Fedus*, P. Fisher,
S. Henderson, A. Kaboth, J. Monroe, T. Sahin*,
G. Sciolla, R. Vanderspek , R. Yamamoto, H. Yegoryan*
Massachusetts Institute of Technology

Note:

* indicates undergraduate students
[1] also MIT
[2] also Harvard University

DM-TPC: detector concept

- Low-pressure CF_4 TPC
  - 50-100 torr→F recoil ~1-2mm
- CF_4 is ideal gas
  - F: spin-dependent interactions
  - Good scintillation efficiency
  - Low transverse diffusion
  - Non flammable, non toxic
- CCD readout
  - Image scintillation photons produced in avalanche
    - $\# \gamma_{\text{scintillation}} \propto \# e_{\text{ionization}}$
  - Low-cost, proven technology
- Amplification region (camera) serves 2 drift regions



Animation by AnaMaria Piso

Detector concept



The amplification region

Original design: wires



- Pros: simplicity, high gains (~10^4-10^5)

What we measure

3 fundamental measurements (in hand)

- E_{recoil} from total scintillation light
- Length of recoil
- Sense of direction (head-tail)
  - Gains an additional order of magnitude
    - A.Green, B.Morgan(astro-ph/0609115)
  - dE/dx <u>decreases</u> along recoil track
    - Low energy, below Bragg peak



Bragg curve for 80 keV F recoil from WIMP in CF_4

Additional features

- PMT provides 3rd coordinate and trigger

No existing experiment had demonstrated head-tail capability

- Our first goal: demonstration of head-tail

First prototype

Drift distance: 2.6cm,
E=580 V/cm
Wire plane: 10x10 cm^2
Anode: 5 mm pitch, 100μm
Ground: 2 mm pitch, 50μm



CCD Camera
Kodak KAF0401 chip
768x512 (9x9mm)
Cooled (-20C)
Photographic lens (55mm)
Finger Lakes Instrumentation



First prototype



5.5 MeV alphas from ^{241}Am source

Alpha track traveling perpendicular to anode wires (vertical)



Bragg curve for 5.5 MeV alphas





- Alphas emitted parallel to anode wires
  - Wire plane oriented at 45 degrees
- Compare measured dE/dx vs range of the track with SRIM simulation
  - Excellent DATA-MC agreement!

Well understood detector

Effect of diffusion on resolution

- Dark Matter recoils ~ 1-2 mm
  - Resolution<< 1mm; diffusion must be contained
- Resolution vs drift distance measured with 4 α sources





$$\sigma[\mu m] = 324 \oplus 36\sqrt{\Delta z}$$

Drift distance	Resolution
1 cm	340 μm
25 cm	670 μm

Light yield calibration with alphas

Photon yield/keV as a function of anode voltage:



Stable operations for gas gain ~ 10^4-10^5

Recoils from low-energy neutrons

- Nuclear recoils by low-energy neutrons mimic Dark Matter
  - DM: F has lower energy but is better aligned with WIMP direction
- Neutron source: 14 MeV neutrons from D-T tube

Fluorine recoil energy



Fluorine recoil angle wrt wires



Blue band = spread in WIMP direction

Neutron beam set





- Anode wires // neutrons
- F recoils in CF_4 at 180 torr:
  E ~ hundred keV, length ~2 mm

NIM A584:327-333,2008

Observation of "head-tail" in F recoils

Wires at 0 deg:







Wires at 180 deg:

Direction of neutrons







NIM A584:327-333,2008

More quantitative results (DT)

We measure skewness of light yield along wire

$$\gamma(x) = \frac{\mu_3}{\mu_2^{3/2}} = \frac{\langle (x - \langle x \rangle)^3 \rangle}{\langle (x - \langle x \rangle)^2 \rangle^{3/2}}$$

γ>0: neutron travels L to R

γ<0: neutron travels R to L





Black dots: wires @ 0 deg
Open circles: wires @180 deg

arXiv:0801.2687

Head-tail with ^{252}Cf source

Very preliminary results (few hours run at P~200 torr)



Observed head-tail for E>200 keV

MC studies: 200 GeV WIMP



Background rejection

- Blind to gammas
  - 8 hours run with 8 μCi ^{137}Cs inside prototype: no evts
  - Rejection factor ~$2/10^6$
- Excellent discrimination against α and e-
  - By measuring both energy and length of recoil
    - For pressure of 50 torr
      - WIMP/neutrons: 30 keV --> 1 mm
      - electrons: 15 keV e- (same ionization) --> 30 mm
      - alphas: 7 keV α --> 1 mm (below threshold)
- Neutrons
  - Passive and active neutron shielding
  - <u>Directionality!</u>

Recent progress: wires --> meshes

- All results shown so far obtained with wire-based detectors
- But wire detectors have serious limitations….





- We recently moved to mesh-based amplification regions

New amplification region: meshes

- Pros: additional coordinate at no additional readout cost
- Cons: more R&D needed

First implementation:



Eventually:



α particles with meshes

V_{anode}=2kV (E~1.5·10^5 V/cm)
0.3mm foil, 200Torr



NB: 1D --> 2D at no additional cost!

^{252}Cf neutrons
100 Torr

Nuclear recoils: very preliminary



~150 keV



~350 keV



~560 keV ← Neutron direction



C: a



or



2nd generation prototype



Drift: ≥ 25mm

Amplification ≥ 5mm

Anode Wires: 7.5mm pitch, 30 μm diameter

Wire Frame: 20x20 cm^2

Image View: ~16 cm diameter

Wire planes from V. Polychronakos (BNL)

CCD Camera: Apogee U2-ME

Kodak KAF-1603ME, 1536x1024 pixels

Lens: Schneider Xenon 0.95/17

First results from 2nd prototype

- Example: deployment of PMT



Po



PMT signal skewed toward left: track tilted toward wire plane

More on PMT signal





Red track from Po: narrow, small tilt away from plane
Blue track not from Po: wide, big tilt wrt wire plane

Next step: ~ m^3 detector

- Goals
  - **Set scientifically competitive limit on spin-dependent interactions with directionality**
  - Prove detector technology on realistic scale
  - Underground backgrounds studies
- Detector
  - Mass: 250-500 g/m^3 for P=50-100 torr
  - 1 year underground run: 90-180 kg-day / m^3
  - Assuming 4 m^3: **350-700 kg-day**
    - SIMPLE/PICASSO 2 Kg-day; CDMS 34(12) Kg-day for Ge(Si) runs
- Timescale
  - Design and R&D: 2007-2008
  - Commissioning and underground run: 2008-2009

Preliminary

Prototype of ~m^3 detector



Preliminary

More on ~ m^3 detector







- 4 amplification regions --> 8 x 25cm drift regions
  - Wires: squared frames; V_{active}~ 2 m^3
  - Triple mesh: circular frames; V_{active}~ 4 m^3
- Low-cost CCD cameras inside active volume (20°C)
- 25-50 cameras/plane; KAI220 chip & 0.95/25 lens

Very preliminary

Sensitivity of 4 m^3 DM-TPC detector



Current best limits on Spin-Dependent interaction

Eventually: full scale detector

- Scale ~100 modules
  - Big for Dark Matter, but small w.r.t. LHC detectors or Super-K
- Simple for today HEP standards
  - E.g.: trigger rates ~ 1 Hz :-)
  - Challenges: purity of materials!
- Mass
  - ~100 kg for P=50-100 torr
  - 3 years of running: ~10^5 Kg day
  - Reach ~ 10^{-10}-10^{-9} cm^2
- Very preliminary cost estimate
  - <~$100K/module
- Shielding
  - Active and passive, a la Braidwood: <$2M

Can we find space for it?

- Large DUSEL cavern at 6,000 mwe is ideal for our needs



From J. Kotcher's presenation at HEPAP meeting on 7/14/2007

Conclusion

- DMTPC collaboration is making rapid progress toward development of new Dark Matter detector
  - Directionality, spin-dependent interactions, optical readout (<$)
- Prototype I proved detector concept (2006-2007)
  - First observation of head-tail effect in low-energy neutrons
    - NIM A584:327-333,2008
- ~1 m^3 module (2007-2009)
  - Design and build (2007-2008): meshes, PMT, better CCD…
  - One year of data taking underground (2008-2009)
    - Competitive limit on spin-dependent cross-section w/directionality
    - Study backgrounds, perfect detector design
- Large DM-TPC detector is an ideal candidate for DUSEL
  - Large cavern at 6,000 mwe ideal for TPC for directional DM detection