

Pseudorapidity Distribution of Charged Particles in $d + \text{Au}$ Collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$

B. B. Back,¹ M. D. Baker,² M. Ballintijn,⁴ D. S. Barton,² B. Becker,² R. R. Betts,⁶ A. A. Bickley,⁷ R. Bindel,⁷ W. Busza,⁴ A. Carroll,² M. P. Decowski,⁴ E. García,⁶ T. Gburek,³ N. George,² K. Gulbrandsen,⁴ S. Gushue,² C. Halliwell,⁶ J. Hamblen,⁸ A. S. Harrington,⁸ C. Henderson,⁴ D. J. Hofman,⁶ R. S. Hollis,⁶ R. Hołyński,³ B. Holzman,² A. Iordanova,⁶ E. Johnson,⁸ J. L. Kane,⁴ N. Khan,⁸ P. Kulinich,⁴ C. M. Kuo,⁵ J. W. Lee,⁴ W. T. Lin,⁵ S. Manly,⁸ A. C. Mignerey,⁷ R. Nouicer,^{2,6} A. Olszewski,³ R. Pak,² I. C. Park,⁸ H. Pernegger,⁴ C. Reed,⁴ C. Roland,⁴ G. Roland,⁴ J. Sagerer,⁶ P. Sarin,⁴ I. Sedykh,² W. Skulski,⁸ C. E. Smith,⁶ P. Steinberg,² G. S. F. Stephens,⁴ A. Sukhanov,² M. B. Tonjes,⁷ A. Trzupek,³ C. Vale,⁴ G. J. van Nieuwenhuizen,⁴ R. Verrier,⁴ G. I. Veres,⁴ F. L. H. Wolfs,⁸ B. Wosiek,³ K. Woźniak,³ B. Wyslouch,⁴ and J. Zhang⁴

(PHOBOS Collaboration)

¹Argonne National Laboratory, Argonne, Illinois 60439-4843, USA

²Brookhaven National Laboratory, Upton, New York 11973-5000, USA

³Institute of Nuclear Physics, Kraków, Poland

⁴Massachusetts Institute of Technology, Cambridge, Massachusetts 02139-4307, USA

⁵National Central University, Chung-Li, Taiwan

⁶University of Illinois at Chicago, Chicago, Illinois 60607-7059, USA

⁷University of Maryland, College Park, Maryland 20742, USA

⁸University of Rochester, Rochester, New York 14627, USA

(Received 10 November 2003; published 16 August 2004)

The measured pseudorapidity distribution of primary charged particles in minimum-bias $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$ is presented for the first time. This distribution falls off less rapidly in the gold direction as compared to the deuteron direction. The average value of the charged particle pseudorapidity density at midrapidity is $\langle dN_{\text{ch}}/d\eta \rangle_{|\eta| \leq 0.6} = 9.4 \pm 0.7(\text{syst})$ and the integrated primary charged particle multiplicity in the measured region is $82 \pm 6(\text{syst})$. Estimates of the total charged particle production, based on extrapolations outside the measured pseudorapidity region, are also presented. The pseudorapidity distribution, normalized to the number of participants in $d + \text{Au}$ collisions, is compared to those of $\text{Au} + \text{Au}$ and $p + \bar{p}$ systems at the same energy. The $d + \text{Au}$ distribution is also compared to the predictions of the parton saturation model, as well as microscopic models.

DOI: 10.1103/PhysRevLett.93.082301

PACS numbers: 25.75.-q

The pseudorapidity distribution of charged particles in $d + \text{Au}$ collisions is important for understanding the evolution of the system created in more complicated $\text{Au} + \text{Au}$ collisions, and may provide a constraint on the initial state parton density. The energy and centrality dependence of pseudorapidity distributions in $\text{Au} + \text{Au}$ collisions measured at BNL Relativistic Heavy Ion Collider (RHIC) [1–3] are consistent with the approach based on the ideas of parton saturation [4,5] and semi-classical QCD [6]. Measurements of $d + \text{Au}$ collisions may be the ideal way to search for the onset of gluon saturation, since the system should be much simpler than the one studied in $\text{Au} + \text{Au}$ collisions. The results of $d + \text{Au}$ collisions are therefore crucial for testing the saturation approach [7] as well as microscopic models (HIJING [8], RQMD [9], and AMPT [10]).

In this Letter we present the first measurement of the minimum-bias pseudorapidity distribution of primary charged particles ($dN_{\text{ch}}/d\eta$) produced in collisions of deuterons with gold nuclei at a nucleon-nucleon center-

of-mass energy, $\sqrt{s_{NN}}$, of 200 GeV. The pseudorapidity, η , is defined as $\eta = -\ln[\tan(\theta/2)]$, where θ is the emission angle relative to the direction of the deuteron beam. The data were obtained with the PHOBOS detector at the Relativistic Heavy Ion Collider at Brookhaven National Laboratory. The data were collected using the multiplicity array [11], covering $|\eta| \leq 5.4$. The array consisted of a barrel of silicon detectors surrounding the beam pipe in the central rapidity region (“Octagon”), and six forward silicon counters, three on each side of the interaction point (“Rings”). The multiplicity array used in $d + \text{Au}$ collisions was the same as that for $\text{Au} + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$ [12]. The detector setup also included two sets of 16 scintillator counters, covering $3 < |\eta| < 4.5$. These counters were used in the primary event trigger and in the offline event selection.

The results presented in this Letter were obtained by two analysis methods: a “hit-counting” method and an “analog” method. The details of the analysis procedure of the two methods leading to the measurement of

$dN_{\text{ch}}/d\eta$ can be found in Ref. [1]. The measured pseudorapidity distribution was corrected for particles which were absorbed or produced in the surrounding material and for feed-down products from weak decays of neutral strange particles. Uncertainties in $dN_{\text{ch}}/d\eta$ associated with these corrections range from 6% in the Octagon and up to 28% in the Rings. These corrections dominate the systematic errors.

Because of the low multiplicity in $d + \text{Au}$ collisions, a new algorithm for collision vertex reconstruction was developed, using the hit position and energy deposited in the Octagon detector. Monte Carlo studies using HIJING [8] and GEANT [13], as well as comparisons with the vertex information from reconstructed tracks, show a resolution of $\sigma_{\text{vtx}}(z) = 1.4$ cm for the most peripheral events and $\sigma_{\text{vtx}}(z) = 0.8$ cm for the most central events, z being the distance along the beam axis.

In order to determine the minimum-bias pseudorapidity distribution for this analysis, it is necessary to correct for the trigger and vertex-finding efficiencies. Since these efficiencies are functions of multiplicity (and therefore centrality), the most model-independent approach is to measure the pseudorapidity distribution in narrow bins of centrality and then integrate over centrality to produce a minimum-bias result. The alternative approach of multiplying by a global Monte Carlo (MC) correction factor (reconstructed/true MC) would rely on our model to get the details of the shape and the centrality dependence right. To further reduce the model dependence, we used five distinct measures for estimating the collision centrality based on the multiplicity (defined by the energy deposited in the silicon detectors) in different regions of pseudorapidity. The first of these centrality measures, E_{Tot} , used data from the full pseudorapidity coverage of PHOBOS, $|\eta| \leq 5.4$. The second measure, E_{Oct} , used data from the more central region $|\eta| \leq 3$. The next two measures, E_{AuDir} and E_{dDir} , used data from the gold direction ($-5.4 \leq \eta \leq -0.5$) and deuteron direction ($0.5 \leq \eta \leq 5.4$), respectively. The final centrality measure, E_{Ring} , used data far from the midrapidity region from the silicon ring counters at $3.0 \leq |\eta| \leq 5.4$.

The centrality measures were calibrated using HIJING and GEANT simulations. In the case of E_{Tot} , E_{Oct} , E_{AuDir} , and E_{dDir} , the distributions in data and MC were very similar once the triggering and event selection were applied to the MC calculation and an additional scaling factor near unity was included to match Monte Carlo simulations and the data distributions in the maximum yield region (highest centrality). This factor (ranging from 0.95 to 1.13) was used in extracting the data centrality bin, but using the 100% efficient Monte Carlo simulation distributions, rather than the biased data distributions. This allowed us to use the zero-bias “true” distribution in the MC to divide the data into equal 10% cross-section bins and to estimate the overall

trigger and vertex-finding efficiency of $\sim 83\%$. This result could be more accurately described as an “inelastic” pseudorapidity density rather than a “minimum-bias” result, as we have attempted to fully correct for the trigger bias. Based on studies of reconstructed and true pseudorapidity distributions from different Monte Carlo simulations, we include a global 5% systematic uncertainty on the pseudorapidity distribution in the reported errors. In the case of E_{Ring} , the MC and data distributions were sufficiently different that the procedure had to be modified. From the calculated efficiencies in each of the




FIG. 1. Minimum-bias pseudorapidity distributions of primary charged particles, $dN_{\text{ch}}/d\eta$, measured for $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV. (a) $dN_{\text{ch}}/d\eta$ distributions obtained by explicit integration over centrality bins for each centrality measure. (b) Final $dN_{\text{ch}}/d\eta$ distribution obtained by averaging over the five minimum-bias distributions. The gray band corresponds to the systematic errors (90% C.L.). The curve corresponds to a triple Gaussian fit to $d + \text{Au}$ collisions.

MC centrality bins, we could derive how many of the events in each true 10% bin would appear in the data. Centrality cuts were then made on the E_{Ring} distribution that gave this efficiency-corrected number of events.

Once the efficiency had been calculated and centrality cuts made, we extracted the average number of participating nucleons $\langle N_{\text{part}} \rangle$ for each centrality bin using HIJING+GEANT. For each of the centrality measures, the minimum-bias pseudorapidity distribution (as well as the estimated number of participants) was obtained by averaging over the ten centrality bins. This averaging was done as the minimum-bias distribution is less sensitive to centrality resolution and multiplicity bias effects than the individual centrality bins. The final distribution was obtained by averaging the five minimum-bias distributions found from the five centrality methods. The total number of participant nucleons, $\langle N_{\text{part}} \rangle$, was estimated to be 8.1 ± 0.7 (syst). To cross check the final distribution obtained by using the silicon centrality measures, a further analysis was performed, only requiring at least one hit in one of the scintillator counter arrays. This analysis was consistent within the present systematic uncertainty.

Figure 1(a) shows the minimum-bias pseudorapidity distributions of primary charged particles measured for $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV obtained with the five centrality measures described above. The distribu-




FIG. 2. $dN_{\text{ch}}/d\eta$ distributions per participant are shown for minimum-bias $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV (solid circles), for central Au + Au collisions (0–6%) [12] (triangles) and for $p + \bar{p}$ collisions from UA5 [19] (open squares) at the same energy. The systematic errors are shown by gray bands. The uncertainties on $\langle N_{\text{part}} \rangle$ for $d + \text{Au}$ and Au + Au collisions have been added in quadrature to the gray bands.

tions agree to within 5% everywhere. Figure 1(b) shows the final result obtained by averaging with equal weights over the five distributions in Fig. 1(a). The systematic errors (90% C.L.) are shown as a gray band. The statistical




FIG. 3. Comparison of the measured minimum-bias pseudorapidity distribution for $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV to model predictions. (a) Comparison to the parton saturation [7] and RQMD [9] models. (b) Comparison to the predictions of the HIJING [8] model and the published AMPT model with and without final-state interactions (FSI) [10]. (c) Quantitative evaluation of the model predictions, expressed as the ratio of the model prediction to the data. The gray band corresponds to the systematic errors (90% C.L.) on the data.

errors are negligible. Figure 1(b) clearly shows that the cross section for particle production is largest in the gold direction ($\eta < 0$), and smallest in the deuteron direction ($\eta > 0$). By fitting the gold side using a Gaussian in the region $-3 \leq \eta \leq -0.5$, the peak centroid was found to be at $\eta \sim -1.9$. This peak position is much closer to midrapidity than predicted by the parton saturation model [7], see Fig. 3(a). It may be interesting to note that a double Gaussian decomposition of the Au + Au distribution at $\sqrt{s_{NN}} = 200$ GeV [12] for the 0–6% most central collisions places the Gaussian centroids at $\eta \sim \pm 1.9$, see Fig. 2. The significance of this observation is, however, not obvious, and may indeed be accidental.

For $d + \text{Au}$ collisions, the measured average value of the charged particle pseudorapidity density at midrapidity is $\langle dN_{\text{ch}}/d\eta \rangle_{|\eta| \leq 0.6} = 9.4 \pm 0.7(\text{syst})$ and the integrated primary charged particle multiplicity in the measured region is $N_{|\eta| \leq 5.4}^{\text{ch}} = 82 \pm 6(\text{syst})$. Since $dN_{\text{ch}}/d\eta$ falls off to only about half of its maximum value at the most negative pseudorapidity of the detector acceptance, it is nontrivial to estimate the total charged particle multiplicity, N_{ch} , for this system. However, using a triple Gaussian fit to the data we obtain the solid curve in Fig. 1(b), which corresponds to a total number of produced charged particles of $N_{\text{tot}}^{\text{ch}} = 99$ and the fit of the upper limit of the systematic errors (upper limit of the gray band) gives $N_{\text{ch}} = 110$. We believe, however, that this represents an extreme upper limit for N_{ch} since experimental data on $p + A$ multiplicity distributions at lower energies [14–18], as well as various theoretical models discussed below, indicate that the falloff at large negative pseudorapidities is more abrupt than indicated by this fit to the data. A more reasonable extrapolation, guided by the expectations based on lower energy $p + A$ data and predictions of the AMPT model, yields an estimate of $N_{\text{ch}} = 5$ in the extrapolated region. Therefore, we estimate that the total charged particle multiplicity in this reaction is $N_{\text{ch}} = 87_{-07}^{+23}$. It should be kept in mind, however, that even this estimate contains a significant contribution from the unmeasured region of pseudorapidity.

Figure 2 shows the normalized minimum-bias $dN_{\text{ch}}/d\eta$ distribution for $d + \text{Au}$ collisions in comparison to the most central (0–6%) Au + Au collisions [12] and non-single-diffractive $p + \bar{p}$ collisions from UA5 [19] at the same energy, $\sqrt{s_{NN}} = 200$ GeV. To be consistent with our previous Au + Au pseudorapidity distribution publications, we have used a normalization factor of $\langle N_{\text{part}}/2 \rangle$ which is unity for $p + \bar{p}$. The distributions from different systems are normalized chiefly to enable a comparison of their shapes. The distribution seen in $d + \text{Au}$ collisions at RHIC has features similar to those seen in $p + A$ collisions at lower energies [14–18]. Compared to $p + \bar{p}$ collisions, there is a significant increase in particle production in the gold fragmentation region and a reduction of particle production at positive pseudorapidity in

the deuteron direction. The overall normalized production of particles is approximately the same as in proton (anti-) proton collisions [19].

Figure 3(a) shows a comparison of the measured minimum-bias pseudorapidity distribution to the published predictions of the parton saturation model [7] and the results from RQMD [9]. It is evident that these models are inconsistent with the data. The parton saturation model overestimates the height of the gold side peak, underestimates its width, and predicts the peak at $\eta \sim -3$ rather than $\eta = -1.9$ as in the data. On the deuteron side, the model underestimates the charged particle production. RQMD drastically underpredicts the particle production and has surprisingly little pseudorapidity asymmetry.

Figure 3(b) presents the comparison of the measured $dN_{\text{ch}}/d\eta$ distribution to the predictions of HIJING [8] and the published AMPT calculation [10]. The HIJING calculation (dash-dotted curve) reproduces the deuteron side and the peak of the gold side, but fails to reproduce the tail in the gold direction ($\eta < -2.5$). The AMPT model uses HIJING for the initial stage of the collision, adding a “popcorn mechanism” for baryon-antibaryon production [10,20] and final-state interactions (FSI). The solid curve is the default, while the dashed curve excludes FSI. We see that both final-state interactions and the popcorn mechanism appear to broaden the gold side peak, leading to a moderate increase of the particle multiplicity in the region $\eta \leq -3.5$. The ratio of the model predictions to data is shown in Fig. 3(c).

In summary, the pseudorapidity distribution of charged particles produced in $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV has been measured. The distribution is rather broad and peaked in the gold direction similar to observations in $p + A$ collisions at lower collision energies. The measured pseudorapidity distribution is compared with the predictions of the parton saturation model as well as microscopic models. While AMPT predictions fall close to the data, the saturation model overestimates the asymmetry between particle production in the gold and deuteron hemispheres.

We thank RHIC Operations for providing the variety of colliding systems. This work was partially supported by U.S. DOE Grants No. DE-AC02-98CH10886, No. DE-FG02-93ER40802, No. DE-FC02-94ER40818, No. DE-FG02-94ER40865, No. DE-FG02-99ER41099, and No. W-31-109-ENG-38, U.S. NSF Grants No. 9603486, No. 9722606, and No. 0072204, Polish KBN Grant No. 2-P03B-10323 and No. NSC of Taiwan Contract No. NSC 89-2112-M-008-024.

-
- [1] B. B. Back *et al.*, Phys. Rev. Lett., **87**, 102303 (2001).
 - [2] B. B. Back *et al.*, Phys. Rev. C **65**, 061901(R) (2002).
 - [3] I. G. Bearden *et al.*, Phys. Rev. Lett. **88**, 202301 (2002).
 - [4] D. Kharzeev *et al.* Phys. Lett. B **523**, 79 (2001).

- [5] L.V. Gribov *et al.*, Phys. Rep. **100**, 1 (1983).
[6] L. McLerran and R. Venugopalan, Phys. Rev. D **49**, 2233 (1994); Phys. Rev. D **50**, 2225 (1994).
[7] D. Kharzeev *et al.*, Nucl. Phys. **A730**, 448 (2004).
[8] M. Gyulassy *et al.*, Comput. Phys. Commun. **83**, 307 (1994). We used HIJING v1.381 with standard parameter settings.
[9] H. Sorge, Phys. Rev. C **52**, 3291 (1995).
[10] Zi-wei Lin *et al.*, Phys. Rev. C **68**, 054904 (2003).
[11] B. B. Back *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **499**, 603 (2003).
[12] B. B. Back *et al.*, Phys. Rev. Lett. **91**, 052303 (2003).
[13] R. Brun *et al.*, GEANT 3.21, CERN Program Library Long Write-up W5013, 1994.
[14] J. E. Elias *et al.*, Phys. Rev. D **22**, 13 (1980).
[15] C. de Marzo *et al.*, Phys. Rev. D **26**, 1019 (1982).
[16] D. H. Brick *et al.*, Phys. Rev. D **41**, 765 (1990).
[17] W. Busza, Acta Phys. Pol. B **8**, 333 (1977).
[18] W. Busza and R. Ledoux, Annu. Rev. Nucl. Part. Sci. **38**, 119 (1988).
[19] G. J. Alner *et al.*, Z. Phys. C **33**, 1 (1986).
[20] Zi-wei Lin *et al.*, Nucl. Phys. **A698**, 375 (2002).